This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

FIGURE 1









p = 0.119

Ισ·


Vehicle


FIGURE 4





Vehicle $\mathcal{I}_{\mathcal{Q}}$










FIGURE 8A, BB








Figure 8C





Figure 8D


FIGURE 9A,9B

A




5

B



10

FIGURE 10



* = p < 0.05 vs. vehicle






FIGURE 12A,12B















FIGURE 14A, 148, 14C, 14 D

Stimulation of the Levels of 3T3-L1 Adipocyte Glut-1



Stimulation of the Levels of 3T3-L1 Adipocyte Glut-4











FIG. 15A


Study-2 (5mg/mouse)				
Day	Animals alive			
	0	3		
	1	3		
	2	3		
	3	3		
	4	3		




Day		mals alive
Duy	٥	2
	4	3
		3
	2	
	3	3
	4	3
		İ
		1











٠.

FIGURE 19











FIGURE 22

Summary of studies of binding of $\pm \alpha$ to nuclear receptors

PPAR	Binding Assay IC ₅₀ (µM)	Cofactor Association EC ₅₀ (µM)	Transactivation EC ₅₀ (μM)
PPAR-α (human)	>50	> 50	No activity
PPAR-a (mouse)		1	No activity
PPAR-y (human)	> 50	>50	No activity
PPAR-6 (human)	× 20	.	i

FIGURE 23



10+ 1+ Vbrt haufin 100 nM 0.5 nM S (MM) 2 E and I stimulate rapid glucose uptake in rat adipocytes Insulfin (n/M) 0.5 . 10 Bust FIGURE 24A,248 20% Lipocril 5 min uptake 14C-Deoxyglucose, sp activity, 315 mCl/mmol É ş Glucose uptake (CPM) (Mn Of) nilusal Gluccae uptales (CPN)




FIGURE 26

Summary of toxicology studies of $\ensuremath{\mathbb{T}} a$

STUDY	STATUS	FINDINGS
Ames mutagenicity (with and without metabolic activation)	complete	No significant toxicity
Acute (single-dose) study, rats	Complete	No significant toxicity
28-Day study, rats Doses: 20, 200, 1000 mg/kg	Complete	No significant toxicity
Cardiovascular study, dogs	Complete	No significant toxicity
Dose range-finding study, dogs Doses: 100, 200, 400, 1000 mg/kg	Complete	No significant toxicity
28-Day study, dogs	Complete	No significant toxicity